

Quarkonium theory

Alexander Rothkopf

Faculty of Science and Technology
Department of Mathematics and Physics
University of Stavanger

References:

S.Kim, P. Petreczky, A.R., JHEP 1811 (2018) 088


P. Petreczky, A.R., J. Weber, NPA982 (2019) 735

D. Lafferty, A.R. arXiv:1906.00035


Y. Akamatsu et. al. JHEP 1807 (2018) 029 & (in preparation)

A challenge to theory

- A wealth of high precision data on both flavors from RHIC and LHC




Bottomonium: a non-equilibrium probe of the full QGP evolution




Charmonium: a partially equilibrated probe, sensitive to the late stages

A challenge to theory

-  A wealth of high precision data on both flavors from RHIC and LHC




Bottomonium: a non-equilibrium probe of the full QGP evolution




Charmonium: a partially equilibrated probe, sensitive to the late stages

-  Goal: provide first principles interpretation to intricate phenomenology

Selected theory goals




Selected theory goals



Properties of equilibrium QQ
Robust in-medium masses
from lattice NRQCD

with S.Kim, P. Petreczky:
JHEP 1811 (2018) 088

Selected theory goals



Properties of equilibrium QQ
Robust in-medium masses
from lattice NRQCD


with S.Kim, P. Petreczky:
JHEP 1811 (2018) 088

In-medium QQ potential
Improved extraction from
realistic lattice QCD

Novel parametrization of $V(R)$
for use in phenomenology

with P. Petreczky, J. Weber: NPA982 (2019) 735
with D. Lafferty arXiv:1906.00035

Selected theory goals



Real-time QQ evol. in local thermal equilibrium

Beyond Schrödinger:
Open-quantum-systems
Lindblad equation

Connecting QQS to
EFT language of potential

with Y.Akamatsu et.al. JHEP 1807 (2018) 029
with S.Kajimoto et.al. PRD97 (2018) 014003
with T. Miura et.al. (in progress)

Properties of equilibrium QQ

Robust in-medium masses
from lattice NRQCD

with S.Kim, P. Petreczky:
JHEP 1811 (2018) 088


In-medium QQ potential


Improved extraction from
realistic lattice QCD

Novel parametrization of $V(R)$
for use in phenomenology

with P. Petreczky, J. Weber: NPA982 (2019) 735
with D. Lafferty arXiv:1906.00035


Heavy quarks in lattice QCD


 Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically




Lattice QCD simulation with $Q\bar{Q}$
still too costly for bottom quarks

Heavy quarks in lattice QCD

 Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically




Lattice QCD simulation with $Q\bar{Q}$
still too costly for bottom quarks




Lattice QCD simulation without $Q\bar{Q}$

Heavy quarks in lattice QCD

- Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically



Lattice QCD simulation with $Q\bar{Q}$
still too costly for bottom quarks




Lattice QCD simulation without $Q\bar{Q}$


- Lattice Non-Relativistic QCD (NRQCD)** well established at $T=0$, applicable at $T>0$
- systematic expansion of QCD action in $1/m_Q a$ Thacker, Lepage Phys.Rev. D43 (1991) 196-208
 - our implementation uses $O(1/(m_Q a)^3)$ and leading order Wilson coefficients

Heavy quarks in lattice QCD

- Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically



Lattice QCD simulation with $Q\bar{Q}$
still too costly for bottom quarks



Lattice QCD simulation without $Q\bar{Q}$

- Lattice Non-Relativistic QCD (NRQCD)** well established at $T=0$, applicable at $T>0$

- systematic expansion of QCD action in $1/m_Q a$ Thacker, Lepage Phys.Rev. D43 (1991) 196-208


- our implementation uses $O(1/(m_Q a)^3)$ and leading order Wilson coefficients

- Realistic $N_f=2+1$ HISQ lattices for the QCD medium by HotQCD** HotQCD PRD85 (2012) 054503, PRD90 (2014) 094503

- $m_\pi=161\text{MeV}$ $T = [140 - 407] \text{ MeV}$ $m_b a = [2.759 - 1.559]$ use adaptive step size to
 $T=0: N_t=32-64$ $T = [140 - 251] \text{ MeV}$ $m_c a = [0.757 - 0.427]$ stabilize NRQCD expansion
 $T>0: 48^3 \times 12$



The direct reconstruction challenge

- **Lattice QCD** simulations are similar to a (very) **imperfect detector**




The direct reconstruction challenge

- **Lattice QCD** simulations are similar to a (very) **imperfect detector**





$$D(\tau) = \int_{-2M_Q}^{\infty} d\omega e^{-\tau\omega} \rho(\omega)$$




The direct reconstruction challenge

- **Lattice QCD** simulations are similar to a (very) **imperfect detector**





$$D(\tau) = \int_{-2M_Q}^{\infty} d\omega e^{-\tau\omega} \rho(\omega)$$




The direct reconstruction challenge

- Lattice QCD simulations are similar to a (very) imperfect detector





$$D(\tau) = \int_{-2M_Q}^{\infty} d\omega e^{-\tau\omega} \rho(\omega)$$




The direct reconstruction challenge

- Lattice QCD simulations are similar to a (very) imperfect detector





$$D(\tau) = \int_{-2M_Q}^{\infty} d\omega e^{-\tau\omega} \rho(\omega)$$




- Extraction of spectra ill-posed **unfolding problem**: here via Bayesian inference

The direct reconstruction challenge

- Lattice QCD simulations are similar to a (very) **imperfect detector**



$$D(\tau) = \int_{-2M_Q}^{\infty} d\omega e^{-\tau\omega} \rho(\omega)$$



- Extraction of spectra ill-posed **unfolding problem**: here via Bayesian inference
- Access to Euclidean time diminishes as T increases - different artifacts as @ $T=0$

Bayesian spectral reconstruction

- Inversion of Laplace transform required – highly ill-posed

$$D(\tau) = \int_{-2m_Q}^{\infty} d\omega e^{-\omega\tau} \rho(\omega)$$

Bayesian spectral reconstruction

- Inversion of Laplace transform required – highly ill-posed

$$D_i = \sum_{l=1}^{N_\omega} \Delta\omega_l e^{-\omega_l \tau_i} \rho_l$$

1. N_ω parameters $\rho_l \gg N_\tau$ datapoints
2. data D_i has finite precision

Bayesian spectral reconstruction

- Inversion of Laplace transform required – highly ill-posed

$$D_i = \sum_{l=1}^{N_\omega} \Delta\omega_l e^{-\omega_l \tau_i} \rho_l$$

1. N_ω parameters $\rho_l \gg N_\tau$ datapoints
2. data D_i has finite precision

- Regularize this task using prior information – Bayes introduces prior $P[\rho|I]=\exp[S]$

M. Jarrell, J. Gubernatis, Physics Reports 269 (3) (1996)

$$P[\rho|D, I] \propto P[D|\rho, I] P[\rho|I] \quad \left. \frac{\delta P[\rho|D, I]}{\delta \rho} \right|_{\rho=\rho^{\text{BR}}} = 0$$

Bayesian spectral reconstruction

- Inversion of Laplace transform required – highly ill-posed

$$D_i = \sum_{l=1}^{N_\omega} \Delta\omega_l e^{-\omega_l \tau_i} \rho_l$$

1. N_ω parameters $\rho_l \gg N_\tau$ datapoints
2. data D_i has finite precision

- Regularize this task using prior information – Bayes introduces prior $P[\rho|I]=\exp[S]$

M. Jarrell, J. Gubernatis, Physics Reports 269 (3) (1996)

$$P[\rho|D, I] \propto P[D|\rho, I] P[\rho|I] \quad \left. \frac{\delta P[\rho|D, I]}{\delta \rho} \right|_{\rho=\rho^{\text{BR}}} = 0$$

- BR prior enforces: ρ positive definite, smoothness of ρ , result independent of units

Y. Burnier, A.R.
PRL 111 (2013) 18, 182003

$$S_{BR} = \alpha \int d\omega \left(1 - \frac{\rho}{m} + \log \left[\frac{\rho}{m} \right] \right)$$

Bayesian spectral reconstruction

- Inversion of Laplace transform required – highly ill-posed

$$D_i = \sum_{l=1}^{N_\omega} \Delta\omega_l e^{-\omega_l \tau_i} \rho_l$$

1. N_ω parameters $\rho_l \gg N_\tau$ datapoints
2. data D_i has finite precision

- Regularize this task using prior information – Bayes introduces prior $P[\rho|I] = \exp[S]$

M. Jarrell, J. Gubernatis, Physics Reports 269 (3) (1996)

$$P[\rho|D, I] \propto P[D|\rho, I] P[\rho|I] \quad \left. \frac{\delta P[\rho|D, I]}{\delta \rho} \right|_{\rho=\rho^{\text{BR}}} = 0$$

- BR prior enforces: ρ positive definite, smoothness of ρ , result independent of units

Y. Burnier, A.R.
PRL 111 (2013) 18, 182003

$$S_{BR} = \alpha \int d\omega \left(1 - \frac{\rho}{m} + \log \left[\frac{\rho}{m} \right] \right)$$

- BR prior: better accuracy in sharp peak structures than MEM or BG but prone to ringing

Bayesian spectral reconstruction

- Inversion of Laplace transform required – highly ill-posed

$$D_i = \sum_{l=1}^{N_\omega} \Delta\omega_l e^{-\omega_l \tau_i} \rho_l$$

1. N_ω parameters $\rho_l \gg N_\tau$ datapoints
2. data D_i has finite precision

- Regularize this task using prior information – Bayes introduces prior $P[\rho|I] = \exp[S]$

M. Jarrell, J. Gubernatis, Physics Reports 269 (3) (1996)

$$P[\rho|D, I] \propto P[D|\rho, I] P[\rho|I] \quad \left. \frac{\delta P[\rho|D, I]}{\delta \rho} \right|_{\rho=\rho^{\text{BR}}} = 0$$

- BR prior enforces: ρ positive definite, smoothness of ρ , result independent of units

Y. Burnier, A.R.
PRL 111 (2013) 18, 182003

$$S_{BR} = \alpha \int d\omega \left(1 - \frac{\rho}{m} + \log \left[\frac{\rho}{m} \right] \right)$$

- BR prior: better accuracy in sharp peak structures than MEM or BG but prone to ringing

C. Fischer, J. Pawłowski,
A.R., C. Welzbacher
PRD98 (2018) 014009

$$S_{BR}^{\text{smooth}} = \alpha \int d\omega \left(\kappa \left(\frac{\partial \rho}{\partial \omega} \right)^2 + 1 - \frac{\rho}{m} + \log \left[\frac{\rho}{m} \right] \right)$$

Bayesian spectral reconstruction

- Inversion of Laplace transform required – highly ill-posed

$$D_i = \sum_{l=1}^{N_\omega} \Delta\omega_l e^{-\omega_l \tau_i} \rho_l$$

1. N_ω parameters $\rho_l \gg N_\tau$ datapoints
2. data D_i has finite precision

- Regularize this task using prior information – Bayes introduces prior $P[\rho|I]=\exp[S]$

M. Jarrell, J. Gubernatis, Physics Reports 269 (3) (1996)

$$P[\rho|D, I] \propto P[D|\rho, I] P[\rho|I] \quad \left. \frac{\delta P[\rho|D, I]}{\delta \rho} \right|_{\rho=\rho^{\text{BR}}} = 0$$

- BR prior enforces: ρ positive definite, smoothness of ρ , result independent of units

Y. Burnier, A.R.
PRL 111 (2013) 18, 182003

$$S_{BR} = \alpha \int d\omega \left(1 - \frac{\rho}{m} + \log \left[\frac{\rho}{m} \right] \right)$$


- BR prior: better accuracy in sharp peak structures than MEM or BG but prone to ringing

C. Fischer, J. Pawłowski,
A.R., C. Welzbacher
PRD98 (2018) 014009

$$S_{BR}^{\text{smooth}} = \alpha \int d\omega \left(\kappa \left(\frac{\partial \rho}{\partial \omega} \right)^2 + 1 - \frac{\rho}{m} + \log \left[\frac{\rho}{m} \right] \right)$$

- Application of different approaches improves understanding of regularization artifacts


NRQCD Euclidean correlators




Non-rel. propagator of
a single heavy quark G

Davies, Thacker Phys.Rev. D45 (1992)

NRQCD Euclidean correlators



NRQCD Euclidean correlators



Non-rel. propagator of
a single heavy quark G



Davies, Thacker Phys.Rev. D45 (1992)

QQ propagator
projected to a certain channel

„correlator of QQ wavefct.

$$D_{J/\psi}(\tau) \triangleq \langle \psi_{J/\psi}(\tau) \psi_{J/\psi}^\dagger(0) \rangle$$

Brambilla et. al. Rev.Mod.Phys. 77 (2005) 1423




Euclidean correlation functions at
 $T=0$ and $T>0$


Correlator ratios ($T>0$ vs $T=0$)

raw lattice NRQCD results

$E_{\text{bind}}(T=0) \sim 1.1 \text{ GeV}$




$E_{\text{bind}}(T=0) \sim 0.6 \text{ GeV}$




Correlator ratios ($T>0$ vs $T=0$)

raw lattice NRQCD results

$E_{\text{bind}}(T=0) \sim 1.1 \text{ GeV}$



$E_{\text{bind}}(T=0) \sim 0.6 \text{ GeV}$




■ In-medium modification hierarchically ordered with vacuum binding energy


Correlator ratios ($T > 0$ vs $T = 0$)

raw lattice NRQCD results



$E_{\text{bind}}(T=0) \sim 1.1 \text{ GeV}$



$E_{\text{bind}}(T=0) \sim 0.6 \text{ GeV}$




Correlator ratio approximated from the lattice potential




- In-medium modification hierarchically ordered with vacuum binding energy
- Upward bend compatible with lower in-medium mass (also seen in previous studies)

Lattice NRQCD spectral functions

$b\bar{b}$ S-wave $T=0$ spectra




$b\bar{b}$ S-wave $T>0$ spectra





Lattice NRQCD spectral functions

$b\bar{b}$ S-wave $T=0$ spectra




$b\bar{b}$ S-wave $T>0$ spectra






 Crucial step: **defining correct $T=0$ baseline** in presence of methods artifacts

Lattice NRQCD spectral functions

$b\bar{b}$ S-wave $T=0$ spectra



$b\bar{b}$ S-wave $T>0$ spectra






- Crucial step: **defining correct $T=0$ baseline** in presence of methods artifacts
- For the first time **consistent negative in medium mass shifts** – ordered by E_{bind}

How to improve in the future?

- Lattice community favorite strategy: more simulations @ smaller lattice spacing?

Mock Tests






- No significant improvement of bound state reconstruction on **finer lattices**

How to improve in the future?

- Lattice community favorite strategy: more simulations @ smaller lattice spacing?

Mock Tests




- No significant improvement of bound state reconstruction on **finer lattices**
- Reached the onset of exponential difficulty: progress needs conceptually new ideas

The real-time interquark potential

Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically

Brambilla et.al. Rev.Mod.Phys. 77 (2005) 1423




$$i\partial_t \langle \psi_s(t) \psi_s(0) \rangle = \left(V^{\text{QCD}}(R) + \mathcal{O}(m_Q^{-1}) + \Theta(R, t) \right) \langle \psi_s(t) \psi_s(0) \rangle$$

The real-time interquark potential

- Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically

Brambilla et.al. Rev.Mod.Phys. 77 (2005) 1423




- $V(R)$ is lowest term in a systematic velocity $v=p/m$ expansion


c.f. potential as interaction kernel in Lipmann Schwinger series in talk by Shuai Liu

The real-time interquark potential

- Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically

 Relativistic $T > 0$
field theory


Brambilla et.al. Rev.Mod.Phys. 77 (2005) 1423


 Quantum
mechanics


$$i\partial_t \langle \psi_s(t) \psi_s(0) \rangle = \left(V^{\text{QCD}}(R) + \mathcal{O}(m_Q^{-1}) + \Theta(R, t) \right) \langle \psi_s(t) \psi_s(0) \rangle$$

- $V(R)$ is lowest term in a systematic velocity $v=p/m$ expansion

 c.f. potential as interaction
kernel in Lipmann Schwinger
series in talk by Shuai Liu


- Matching to underlying QCD in the infinite mass limit: Wilson loop

$$\langle \psi_S(R, t) \psi_S^*(R, 0) \rangle_{\text{pNRQCD}} \equiv W_\square(R, t) = \left\langle \text{Tr} \left[\exp \left(-ig \int_\square dx^\mu A_\mu(x) \right) \right] \right\rangle_{\text{QCD}}$$




The real-time interquark potential

- Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically

 Relativistic $T > 0$
field theory


Brambilla et.al. Rev.Mod.Phys. 77 (2005) 1423



 Quantum
mechanics


$$i\partial_t \langle \psi_S(t) \psi_S(0) \rangle = \left(V^{\text{QCD}}(R) + \mathcal{O}(m_Q^{-1}) + \Theta(R, t) \right) \langle \psi_S(t) \psi_S(0) \rangle$$

- $V(R)$ is lowest term in a systematic velocity $v=p/m$ expansion c.f. potential as interaction kernel in Lipmann Schwinger series in talk by Shuai Liu
- Matching to underlying QCD in the infinite mass limit: Wilson loop

$$\langle \psi_S(R, t) \psi_S^*(R, 0) \rangle_{\text{pNRQCD}} \equiv W_\square(R, t) = \left\langle \text{Tr} \left[\exp \left(-ig \int_\square dx^\mu A_\mu(x) \right) \right] \right\rangle_{\text{QCD}}$$


- Wilson loop: potential emerges at late times

$$V(R) = \lim_{t \rightarrow \infty} \frac{i\partial_t W_\square(R, t)}{W_\square(R, t)}$$




The real-time interquark potential

- Exploit $\frac{T}{m_Q} \ll 1, \frac{\Lambda_{\text{QCD}}}{m_Q} \ll 1$ to treat heavy quarks non-relativistically

 Relativistic $T > 0$
field theory


Brambilla et.al. Rev.Mod.Phys. 77 (2005) 1423



 Quantum
mechanics

$$i\partial_t \langle \psi_S(t) \psi_S(0) \rangle = \left(V^{\text{QCD}}(R) + \mathcal{O}(m_Q^{-1}) + \Theta(R, t) \right) \langle \psi_S(t) \psi_S(0) \rangle$$


- $V(R)$ is lowest term in a systematic velocity $v=p/m$ expansion c.f. potential as interaction kernel in Lipmann Schwinger series in talk by Shuai Liu
- Matching to underlying QCD in the infinite mass limit: Wilson loop

$$\langle \psi_S(R, t) \psi_S^*(R, 0) \rangle_{\text{pNRQCD}} \equiv W_\square(R, t) = \left\langle \text{Tr} \left[\exp \left(-ig \int_\square dx^\mu A_\mu(x) \right) \right] \right\rangle_{\text{QCD}}$$

- Wilson loop: potential emerges at late times

$$V(R) = \lim_{t \rightarrow \infty} \frac{i\partial_t W_\square(R, t)}{W_\square(R, t)} \in \mathbb{C}$$

Im[V]: Laine et al. JHEP03 (2007) 054;
Beraudo et. al. NPA 806:312,2008
Brambilla et.al. PRD 78 (2008) 014017



Non-perturbative evaluation of $V(R)$

- How to connect to the Euclidean domain: **spectral functions**

A.R., T.Hatsuda & S.Sasaki
PRL 108 (2012) 162001

$$W_{\square}(R, t) = \int_{-\infty}^{\infty} d\omega e^{-i\omega t} \rho_{\square}(R, \omega) \quad \longleftrightarrow \quad W_{\square}(R, \tau) = \int_{-\infty}^{\infty} d\omega e^{-\omega \tau} \rho_{\square}(R, \omega)$$

Non-perturbative evaluation of $V(R)$

- How to connect to the Euclidean domain: **spectral functions**

A.R., T.Hatsuda & S.Sasaki
PRL 108 (2012) 162001

$$W_{\square}(R, t) = \int_{-\infty}^{\infty} d\omega e^{-i\omega t} \rho_{\square}(R, \omega) \quad \longleftrightarrow \quad W_{\square}(R, \tau) = \int_{-\infty}^{\infty} d\omega e^{-\omega \tau} \rho_{\square}(R, \omega)$$

Spectral Decomposition

$$V^{\text{QCD}}(R) = \lim_{t \rightarrow \infty} \frac{\int_{-\infty}^{\infty} d\omega \omega e^{-i\omega t} \rho_{\square}(R, \omega)}{\int_{-\infty}^{\infty} d\omega e^{-i\omega t} \rho_{\square}(R, \omega)}$$

A.R., T.Hatsuda & S.Sasaki PoS LAT2009 (2009) 162

Non-perturbative evaluation of $V(R)$

- How to connect to the Euclidean domain: **spectral functions**


A.R., T.Hatsuda & S.Sasaki
PRL 108 (2012) 162001

$$W_{\square}(R, t) = \int_{-\infty}^{\infty} d\omega e^{-i\omega t} \rho_{\square}(R, \omega) \quad \longleftrightarrow \quad W_{\square}(R, \tau) = \int_{-\infty}^{\infty} d\omega e^{-\omega \tau} \rho_{\square}(R, \omega)$$

Spectral Decomposition

$$V^{\text{QCD}}(R) = \lim_{t \rightarrow \infty} \frac{\int_{-\infty}^{\infty} d\omega \omega e^{-i\omega t} \rho_{\square}(R, \omega)}{\int_{-\infty}^{\infty} d\omega e^{-i\omega t} \rho_{\square}(R, \omega)}$$

A.R., T.Hatsuda & S.Sasaki PoS LAT2009 (2009) 162



well defined $V(R)$
if low lying Breit-
Wigner present in
Wilson loop
spectral function

For technical details see
Y.B., A.R. PRD86 (2012) 051503

$$V(R) = \omega_0(R) - i\Gamma_0(R)$$

Non-perturbative evaluation of $V(R)$

- How to connect to the Euclidean domain: **spectral functions**


A.R., T.Hatsuda & S.Sasaki
PRL 108 (2012) 162001

$$W_{\square}(R, t) = \int_{-\infty}^{\infty} d\omega e^{-i\omega t} \rho_{\square}(R, \omega) \quad \longleftrightarrow \quad W_{\square}(R, \tau) = \int_{-\infty}^{\infty} d\omega e^{-\omega \tau} \rho_{\square}(R, \omega)$$

Spectral Decomposition

$$V^{QCD}(R) = \lim_{t \rightarrow \infty} \frac{\int_{-\infty}^{\infty} d\omega \omega e^{-i\omega t} \rho_{\square}(R, \omega)}{\int_{-\infty}^{\infty} d\omega e^{-i\omega t} \rho_{\square}(R, \omega)}$$

A.R., T.Hatsuda & S.Sasaki PoS LAT2009 (2009) 162



well defined $V(R)$
if low lying Breit-
Wigner present in
Wilson loop
spectral function
For technical details see
Y.B., A.R. PRD86 (2012) 051503

$$V(R) = \omega_0(R) - i\Gamma_0(R)$$

Spectral Reconstruction

- In case of usual $\Delta W/W=10^{-2}$ statistical uncertainty in W_{\square} : **Bayesian inference**

incorporate prior information to regularize the inversion task (BR method)

- In case of **small $\Delta W/W < 10^{-3}$** statistical uncertainty in W_{\square} also **Pade approximation**

exploit the analyticity of the
Wilson correlator to extract spectra


Latest results on the lattice potential




- Lattices with dynamical u,d,s quarks (HISQ action, HotQCD & TUMQCD)


A. Bazavov et.al. PRD97 (2018) 014510, HotQCD PRD90 (2014) 094503


- realistic $m_\pi \sim 161 \text{ MeV}$ ($T=151-1451 \text{ MeV}$)
- fixed box ($N_s=48$ - **$N_t=12$, $N_t=16$**) & very **high statistics** 4000-9000 realizations
- Pade based extraction for $\text{Re}[V]$ possible

Latest results on the lattice potential

-  Lattices with dynamical u,d,s quarks (HISQ action, HotQCD & TUMQCD)

A. Bazavov et.al. PRD97 (2018) 014510, HotQCD PRD90 (2014) 094503
-  realistic $m_\pi \sim 161 \text{ MeV}$ ($T=151\text{-}1451 \text{ MeV}$)
-  fixed box ($N_s=48 - N_t=12, N_t=16$) & very **high statistics** 4000-9000 realizations
-  Pade based extraction for $\text{Re}[V]$ possible





-  Smooth transition from Cornell @ $T=0$ to Debye screened @ $T>T_c$

Latest results on the lattice potential

- Lattices with dynamical u,d,s quarks (HISQ action, HotQCD & TUMQCD)

A. Bazavov et.al. PRD97 (2018) 014510, HotQCD PRD90 (2014) 094503


- realistic $m_\pi \sim 161 \text{ MeV}$ ($T=151\text{-}1451 \text{ MeV}$)
- fixed box ($N_s=48 - N_t=12, N_\tau=16$) & very **high statistics** 4000-9000 realizations
- Pade based extraction for $\text{Re}[V]$ possible



- Smooth transition from Cornell @ $T=0$ to Debye screened @ $T>T_c$
- Finite $\text{Im}[V]$ above T_c present

An improved Gauss law approach

- For use in phenomenology applications: analytic expression for $\text{Re}[V]$ and $\text{Im}[V]$




$$V_{Q\bar{Q}}^{T=0}(R) = V_C(R) + V_S(R) = -\frac{\alpha_s}{r} + \sigma r + c$$

Strategy:

α_s, σ and c are vacuum prop.
and do not change with T

An improved Gauss law approach

- For use in phenomenology applications: analytic expression for $\text{Re}[V]$ and $\text{Im}[V]$



$$V_{Q\bar{Q}}^{T=0}(R) = V_C(R) + V_S(R) = -\frac{\alpha_s}{r} + \sigma r + c$$

$$\mathcal{G}_a[V(R)] = \vec{\nabla} \left(\frac{\vec{\nabla} V(R)}{R^{a+1}} \right) = -4\pi q \delta^{(3)}(\vec{R})$$

Strategy:

α_s, σ and c are vacuum prop. and do not change with T

$$V(R) = aqR^a$$

Coulombic: $a=-1$ $q=\alpha_s$

$$\vec{\nabla} \left(\vec{\nabla} V_C(R) \right) = -4\pi\alpha_s \delta(\vec{R})$$


String-like: $a=+1$ $q=\sigma$

$$\vec{\nabla} \left(\frac{\vec{\nabla} V_S(R)}{R^2} \right) = -4\pi\sigma \delta(\vec{R})$$

V. V. Dixit,
Mod. Phys. Lett. A 5, 227 (1990)

An improved Gauss law approach

- For use in phenomenology applications: analytic expression for $\text{Re}[V]$ and $\text{Im}[V]$



$$V_{Q\bar{Q}}^{T=0}(R) = V_C(R) + V_S(R) = -\frac{\alpha_S}{r} + \sigma r + c$$

$$\mathcal{G}_a[V(R)] = \vec{\nabla} \left(\frac{\vec{\nabla} V(R)}{R^{a+1}} \right) = -4\pi q \delta^{(3)}(\vec{R})$$

Strategy:

α_s, σ and c are vacuum prop. and do not change with T

$$V(R) = aqR^a$$

Coulombic: $a=-1$ $q=\alpha_s$

$$\vec{\nabla} \left(\vec{\nabla} V_C(R) \right) = -4\pi \alpha_s \delta(\vec{R})$$


String-like: $a=+1$ $q=\sigma$

$$\vec{\nabla} \left(\frac{\vec{\nabla} V_S(R)}{R^2} \right) = -4\pi \sigma \delta(\vec{R})$$

V. V. Dixit,
Mod. Phys. Lett. A 5, 227 (1990)

- Immerse non-perturbative charge in weak coupling HTL medium: permittivity ϵ


original idea: Y.Burnier, A.R. Phys.Lett. B753 (2016) 232 improved derivation D.Lafferty and A.R. arXiv:1906.00035



$$V^{med}(\mathbf{p}) = V^{vac}(\mathbf{p})/\epsilon(\mathbf{p}) \quad \epsilon^{-1}(\vec{p}, m_D) = \frac{p^2}{p^2 + m_D^2} - i\pi T \frac{pm_D^2}{(p^2 + m_D^2)^2}$$

An improved Gauss law approach

- For use in phenomenology applications: analytic expression for $\text{Re}[V]$ and $\text{Im}[V]$



$$V_{Q\bar{Q}}^{T=0}(R) = V_C(R) + V_S(R) = -\frac{\alpha_S}{r} + \sigma r + c$$

$$\mathcal{G}_a[V(R)] = \vec{\nabla} \left(\frac{\vec{\nabla} V(R)}{R^{a+1}} \right) = -4\pi q \delta^{(3)}(\vec{R})$$

Strategy:

α_S, σ and c are vacuum prop. and do not change with T

$$V(R) = aqR^a$$

Coulombic: $a=-1$ $q=\alpha_S$

$$\vec{\nabla} \left(\vec{\nabla} V_C(R) \right) = -4\pi \alpha_S \delta(\vec{R})$$


String-like: $a=+1$ $q=\sigma$

$$\vec{\nabla} \left(\frac{\vec{\nabla} V_S(R)}{R^2} \right) = -4\pi \sigma \delta(\vec{R})$$

V. V. Dixit,
Mod. Phys. Lett. A 5, 227 (1990)

- Immerse non-perturbative charge in weak coupling HTL medium: permittivity ϵ

original idea: Y.Burnier, A.R. Phys.Lett. B753 (2016) 232 improved derivation D.Lafferty and A.R. arXiv:1906.00035




$$V^{med}(\mathbf{p}) = V^{vac}(\mathbf{p})/\epsilon(\mathbf{p}) \quad \epsilon^{-1}(\vec{p}, m_D) = \frac{p^2}{p^2 + m_D^2} - i\pi T \frac{pm_D^2}{(p^2 + m_D^2)^2}$$

$$\mathcal{G}_a[V^{med}(\mathbf{r})] = \mathcal{G}_a \int d^3y (V^{vac}(\mathbf{r} - \mathbf{y}) \epsilon^{-1}(\mathbf{y})) = 4\pi q \epsilon^{-1}(\mathbf{r}, m_D)$$

An improved Gauss law approach

- For use in phenomenology applications: analytic expression for $\text{Re}[V]$ and $\text{Im}[V]$



$$V_{Q\bar{Q}}^{T=0}(R) = V_C(R) + V_S(R) = -\frac{\alpha_S}{r} + \sigma r + c$$

$$\mathcal{G}_a[V(R)] = \vec{\nabla} \left(\frac{\vec{\nabla} V(R)}{R^{a+1}} \right) = -4\pi q \delta^{(3)}(\vec{R})$$

Strategy:

α_s, σ and c are vacuum prop. and do not change with T

$$V(R) = aqR^a$$

Coulombic: $a=-1$ $q=\alpha_s$

$$\vec{\nabla}(\vec{\nabla} V_C(R)) = -4\pi\alpha_s\delta(\vec{R})$$


String-like: $a=+1$ $q=\sigma$

$$\vec{\nabla} \left(\frac{\vec{\nabla} V_S(R)}{R^2} \right) = -4\pi\sigma\delta(\vec{R})$$

V. V. Dixit,
Mod. Phys. Lett. A 5, 227 (1990)

- Immerse non-perturbative charge in weak coupling HTL medium: permittivity ϵ

original idea: Y.Burnier, A.R. Phys.Lett. B753 (2016) 232 improved derivation D.Lafferty and A.R. arXiv:1906.00035



$$V^{med}(\mathbf{p}) = V^{vac}(\mathbf{p})/\epsilon(\mathbf{p}) \quad \epsilon^{-1}(\vec{p}, m_D) = \frac{p^2}{p^2 + m_D^2} - i\pi T \frac{pm_D^2}{(p^2 + m_D^2)^2}$$



$$\mathcal{G}_a[V^{med}(\mathbf{r})] = \mathcal{G}_a \int d^3y (V^{vac}(\mathbf{r} - \mathbf{y})\epsilon^{-1}(\mathbf{y})) = 4\pi q\epsilon^{-1}(\mathbf{r}, m_D)$$

- 3 vacuum parameters and 1 temperature dependent m_D fix both $\text{Re}[V]$ and $\text{Im}[V]$.

Gauss-law solution to $\text{Re}[V]$ & $\text{Im}[V]$

■ Gauss-Law result allows to fit $\text{Re}[V]$ data even in the non-perturbative regime



D.Lafferty and A.R. arXiv:1906.00035




Gauss-law solution to $\text{Re}[V]$ & $\text{Im}[V]$

- Gauss-Law result allows to fit $\text{Re}[V]$ data even in the non-perturbative regime

D.Lafferty and A.R. arXiv:1906.00035





- m_D defined from $\text{Re}[V]$ allows to compute Gauss law prediction for $\text{Im}[V]$




Gauss-law solution to $\text{Re}[V]$ & $\text{Im}[V]$

- Gauss-Law result allows to fit $\text{Re}[V]$ data even in the non-perturbative regime

D.Lafferty and A.R. arXiv:1906.00035




- m_D defined from $\text{Re}[V]$ allows to compute Gauss law prediction for $\text{Im}[V]$
- recently extend the Gauss law to model quarkonium at finite velocity & μ_B



The open quantum systems picture

- Need a general approach to describe quarkonium coupled to a thermal medium
- Overall system is closed, hermitean Hamiltonian: von Neumann equation


$$H = H_{Q\bar{Q}} \otimes I_{med} + I_{Q\bar{Q}} \otimes H_{med} + H_{int} \quad \frac{d\rho}{dt} = -i[H, \rho]$$



The open quantum systems picture

- Need a general approach to describe quarkonium coupled to a thermal medium
- Overall system is closed, hermitean Hamiltonian: von Neumann equation

$$H = H_{Q\bar{Q}} \otimes I_{med} + I_{Q\bar{Q}} \otimes H_{med} + H_{int} \quad \frac{d\rho}{dt} = -i[H, \rho]$$




- Dynamics of the reduced QQbar system: $\rho_{Q\bar{Q}} = \text{Tr}_{med}[\rho]$ $\frac{d\rho_{Q\bar{Q}}}{dt} = ?$
for an EFT result see Brambilla et.al. PRD96 (2017) , 034021

The open quantum systems picture

- Need a general approach to describe quarkonium coupled to a thermal medium
- Overall system is closed, hermitean Hamiltonian: von Neumann equation

$$H = H_{Q\bar{Q}} \otimes I_{med} + I_{Q\bar{Q}} \otimes H_{med} + H_{int} \quad \frac{d\rho}{dt} = -i[H, \rho]$$




- Dynamics of the reduced QQbar system: $\rho_{Q\bar{Q}} = \text{Tr}_{med}[\rho]$ $\frac{d\rho_{Q\bar{Q}}}{dt} = ?$
for an EFT result see Brambilla et.al. PRD96 (2017) , 034021

- Very versatile: time scale hierarchies allow to derive simplified description see Xiaojun Yao's talk on Boltzmann eq.

The open quantum systems picture

- Need a general approach to describe quarkonium coupled to a thermal medium
- Overall system is closed, hermitean Hamiltonian: von Neumann equation

$$H = H_{Q\bar{Q}} \otimes I_{med} + I_{Q\bar{Q}} \otimes H_{med} + H_{int} \quad \frac{d\rho}{dt} = -i[H, \rho]$$



- Dynamics of the reduced QQbar system: $\rho_{Q\bar{Q}} = \text{Tr}_{med}[\rho]$ $\frac{d\rho_{Q\bar{Q}}}{dt} = ?$
for an EFT result see Brambilla et.al. PRD96 (2017) , 034021

- Very versatile: time scale hierarchies allow to derive simplified description see Xiaojun Yao's talk on Boltzmann eq.


- Derivation via path integral formalism: **Feynman-Vernon influence functional**
for details see Y. Akamatsu, Phys.Rev. D87 (2013) 4, 045016 and arXiv:1403.5783

$$\rho(t, x, y, X, Y) = \int dx_0 dy_0 dX_0 dY_0 \rho(0, x_0, y_0, X_0, Y_0) \int_{x_0, y_0, X_0, Y_0}^{x, y, X, Y} \mathcal{D}[\bar{x}, \bar{y}, \bar{X}, \bar{Y}] e^{iS[\bar{x}, \bar{X}] - iS[\bar{y}, \bar{Y}]}$$

The open quantum systems picture

- Need a general approach to describe quarkonium coupled to a thermal medium
- Overall system is closed, hermitean Hamiltonian: von Neumann equation

$$H = H_{Q\bar{Q}} \otimes I_{med} + I_{Q\bar{Q}} \otimes H_{med} + H_{int} \quad \frac{d\rho}{dt} = -i[H, \rho]$$



- Dynamics of the reduced QQbar system: $\rho_{Q\bar{Q}} = \text{Tr}_{med}[\rho]$ $\frac{d\rho_{Q\bar{Q}}}{dt} = ?$
for an EFT result see Brambilla et.al. PRD96 (2017) , 034021
- Very versatile: time scale hierarchies allow to derive simplified description see Xiaojun Yao's talk on Boltzmann eq.
- Derivation via path integral formalism: **Feynman-Vernon influence functional**
for details see Y. Akamatsu, Phys.Rev. D87 (2013) 4, 045016 and arXiv:1403.5783

$$\rho(t, x, y, X, Y) = \int dx_0 dy_0 dX_0 dY_0 \rho(0, x_0, y_0, X_0, Y_0) \int_{x_0, y_0, X_0, Y_0}^{x, y, X, Y} \mathcal{D}[\bar{x}, \bar{y}, \bar{X}, \bar{Y}] e^{iS[\bar{x}, \bar{X}] - iS[\bar{y}, \bar{Y}]}$$


$$\rho_{Q\bar{Q}}(t, x, y) = \int dX dY \delta(X - Y) \rho(t, x, y, X, Y)$$

The open quantum systems picture

- Need a general approach to describe quarkonium coupled to a thermal medium

- Overall system is closed, hermitean Hamiltonian: von Neumann equation

$$H = H_{Q\bar{Q}} \otimes I_{med} + I_{Q\bar{Q}} \otimes H_{med} + H_{int} \quad \frac{d\rho}{dt} = -i[H, \rho]$$



- Dynamics of the reduced QQbar system: $\rho_{Q\bar{Q}} = \text{Tr}_{med}[\rho]$ $\frac{d\rho_{Q\bar{Q}}}{dt} = ?$
for an EFT result see Brambilla et.al. PRD96 (2017) , 034021

- Very versatile: time scale hierarchies allow to derive simplified description see Xiaojun Yao's talk on Boltzmann eq.

- Derivation via path integral formalism: **Feynman-Vernon influence functional**
for details see Y. Akamatsu, Phys.Rev. D87 (2013) 4, 045016 and arXiv:1403.5783


$$\rho(t, x, y, X, Y) = \int dx_0 dy_0 dX_0 dY_0 \rho(0, x_0, y_0, X_0, Y_0) \int_{x_0, y_0, X_0, Y_0}^{x, y, X, Y} \mathcal{D}[\bar{x}, \bar{y}, \bar{X}, \bar{Y}] e^{iS[\bar{x}, \bar{X}] - iS[\bar{y}, \bar{Y}]}$$

$$\rho_{Q\bar{Q}}(t, x, y) = \int dx_0 dy_0 \rho_{Q\bar{Q}}(0, x, y) \int_{x_0, y_0}^{x, y} \mathcal{D}[\bar{x}, \bar{y}] e^{iS_{Q\bar{Q}}[\bar{x}] - iS_{Q\bar{Q}}[\bar{y}] + iS_{FV}[\bar{x}, \bar{y}]}$$

The open quantum systems picture

- Need a general approach to describe quarkonium coupled to a thermal medium
- Overall system is closed, hermitean Hamiltonian: von Neumann equation

$$H = H_{Q\bar{Q}} \otimes I_{med} + I_{Q\bar{Q}} \otimes H_{med} + H_{int} \quad \frac{d\rho}{dt} = -i[H, \rho]$$



- Dynamics of the reduced QQbar system: $\rho_{Q\bar{Q}} = \text{Tr}_{med}[\rho]$ $\frac{d\rho_{Q\bar{Q}}}{dt} = ?$
for an EFT result see Brambilla et.al. PRD96 (2017) , 034021

- Very versatile: time scale hierarchies allow to derive simplified description see Xiaojun Yao's talk on Boltzmann eq.

- Derivation via path integral formalism: **Feynman-Vernon influence functional**
for details see Y. Akamatsu, Phys.Rev. D87 (2013) 4, 045016 and arXiv:1403.5783

$$\rho(t, x, y, X, Y) = \int dx_0 dy_0 dX_0 dY_0 \rho(0, x_0, y_0, X_0, Y_0) \int_{x_0, y_0, X_0, Y_0}^{x, y, X, Y} \mathcal{D}[\bar{x}, \bar{y}, \bar{X}, \bar{Y}] e^{iS[\bar{x}, \bar{X}] - iS[\bar{y}, \bar{Y}]}$$

$$\rho_{Q\bar{Q}}(t, x, y) = \int dx_0 dy_0 \rho_{Q\bar{Q}}(0, x, y) \int_{x_0, y_0}^{x, y} \mathcal{D}[\bar{x}, \bar{y}] e^{iS_{Q\bar{Q}}[\bar{x}] - iS_{Q\bar{Q}}[\bar{y}] + \boxed{iS_{FV}[\bar{x}, \bar{y}]}}$$

medium - QQ interaction

The Lindblad equation

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

The Lindblad equation

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

The Lindblad equation

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- At the moment model simply uses 0th order approximation $S_{pot}[Re[V] - i Im[V]]$

The Lindblad equation

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation


$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- At the moment model simply uses 0th order approximation $S_{pot}[Re[V] - i Im[V]]$

Krouppa et. al., PRDD97 (2018) 016017
[STAR Collaboration] Z. Ye
Nucl.Part.Phys.Proc. 289-290 (2017) 401



The Lindblad equation

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation


$$S_{FV} \approx S_{\text{pot}}[Re[V]] + S_{\text{fluct}}[Im[V]] + S_{\text{diss}}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation


$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- At the moment model simply uses 0th order approximation $S_{\text{pot}}[Re[V] - i Im[V]]$

Kruppa et. al., PRDD97 (2018) 016017
[STAR Collaboration] Z. Ye
Nucl.Part.Phys.Proc. 289-290 (2017) 401



Comparison to 1st order approximation



S.Kajimoto, Y.Akamatsu, M. Asakawa,
A.R., PRD97 (2018) 014003

Quantum state diffusion

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- Actually: unravel into wavefunction stochastic dynamics: **Quantum State Diffusion**
T. Miura, Y. Akamatsu, M. Asakawa, S. Kajimoto, A.R., in preparation

$$\begin{aligned} |d\psi\rangle &= |\psi(t+dt)\rangle - |\psi(t)\rangle \\ &= -iH|\psi(t)\rangle dt + \sum_n \begin{pmatrix} 2\langle L_n^\dagger \rangle_\psi L_n - L_n^\dagger L_n \\ -\langle L_n^\dagger \rangle_\psi \langle L_n \rangle_\psi \end{pmatrix} |\psi(t)\rangle dt \\ &\quad + \sum_n (L_n - \langle L_n \rangle_\psi) |\psi(t)\rangle d\xi_n, \end{aligned}$$

$$\begin{aligned} L_{\vec{k},a} &= \sqrt{\frac{D(\vec{k})}{2}} \left[1 - \frac{\vec{k}}{4MT} \cdot \left(\frac{1}{2} \vec{P}_{CM} + \hat{p} \right) \right] e^{i\vec{k} \cdot \hat{r}/2} (t^a \otimes 1) \\ &\quad - \sqrt{\frac{D(\vec{k})}{2}} \left[1 - \frac{\vec{k}}{4MT} \cdot \left(\frac{1}{2} \vec{P}_{CM} - \hat{p} \right) \right] e^{-i\vec{k} \cdot \hat{r}/2} (1 \otimes t^{a*}) \end{aligned}$$

$$\tilde{D}(k) = g^2 T \frac{\pi m_D^2}{k(k^2 + m_D^2)^2}, \quad m_D = gT \sqrt{\frac{N_c}{3} + \frac{N_f}{6}}$$

high temperature weak coupling parameters

Quantum state diffusion

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- Actually: unravel into wavefunction stochastic dynamics: **Quantum State Diffusion**

T. Miura, Y. Akamatsu, M. Asakawa, S. Kajimoto, A.R., in preparation

$$\begin{aligned} |d\psi\rangle &= |\psi(t+dt)\rangle - |\psi(t)\rangle \\ &= -iH|\psi(t)\rangle dt + \sum_n \begin{pmatrix} 2\langle L_n^\dagger \rangle_\psi L_n - L_n^\dagger L_n \\ -\langle L_n^\dagger \rangle_\psi \langle L_n \rangle_\psi \end{pmatrix} |\psi(t)\rangle dt \\ &\quad + \sum_n (L_n - \langle L_n \rangle_\psi) |\psi(t)\rangle d\xi_n, \end{aligned}$$

$$\begin{aligned} L_{\vec{k},a} &= \sqrt{\frac{D(\vec{k})}{2}} \left[1 - \frac{\vec{k}}{4MT} \cdot \left(\frac{1}{2} \vec{P}_{CM} + \hat{p} \right) \right] e^{i\vec{k} \cdot \hat{r}/2} (t^a \otimes 1) \\ &\quad - \sqrt{\frac{D(\vec{k})}{2}} \left[1 - \frac{\vec{k}}{4MT} \cdot \left(\frac{1}{2} \vec{P}_{CM} - \hat{p} \right) \right] e^{-i\vec{k} \cdot \hat{r}/2} (1 \otimes t^{a*}) \end{aligned}$$

$$\tilde{D}(k) = g^2 T \frac{\pi m_D^2}{k(k^2 + m_D^2)^2}, \quad m_D = gT \sqrt{\frac{N_c}{3} + \frac{N_f}{6}}$$

high temperature weak coupling parameters

- First “derivation” of phenomenological models based on nonlinear Schrödinger equation

c.f. e.g. R. Katz, P. Gossiaux Annals Phys. 368 (2016) 267

Quantum state diffusion

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- Actually: unravel into wavefunction stochastic dynamics: **Quantum State Diffusion**

T. Miura, Y. Akamatsu, M. Asakawa, S. Kajimoto, A.R., in preparation

$$\begin{aligned} |d\psi\rangle &= |\psi(t+dt)\rangle - |\psi(t)\rangle \\ &= -iH|\psi(t)\rangle dt + \sum_n \begin{pmatrix} 2\langle L_n^\dagger \rangle_\psi L_n - L_n^\dagger L_n \\ -\langle L_n^\dagger \rangle_\psi \langle L_n \rangle_\psi \end{pmatrix} |\psi(t)\rangle dt \\ &\quad + \sum_n (L_n - \langle L_n \rangle_\psi) |\psi(t)\rangle d\xi_n, \end{aligned}$$

$$\begin{aligned} L_{\vec{k},a} &= \sqrt{\frac{D(\vec{k})}{2}} \left[1 - \frac{\vec{k}}{4MT} \cdot \left(\frac{1}{2} \vec{P}_{CM} + \hat{p} \right) \right] e^{i\vec{k} \cdot \hat{r}/2} (t^a \otimes 1) \\ &\quad - \sqrt{\frac{D(\vec{k})}{2}} \left[1 - \frac{\vec{k}}{4MT} \cdot \left(\frac{1}{2} \vec{P}_{CM} - \hat{p} \right) \right] e^{-i\vec{k} \cdot \hat{r}/2} (1 \otimes t^{a*}) \end{aligned}$$

$$\tilde{D}(k) = g^2 T \frac{\pi m_D^2}{k(k^2 + m_D^2)^2}, \quad m_D = gT \sqrt{\frac{N_c}{3} + \frac{N_f}{6}}$$

high temperature weak coupling parameters

- First “derivation” of phenomenological models based on nonlinear Schrödinger equation

c.f. e.g. R. Katz, P. Gossiaux Annals Phys. 368 (2016) 267

- First genuine Lindblad implementation: previous works could not maintain positivity of ρ

D. De Boni, JHEP 1708 (2017) 064

Quantum state diffusion

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$



- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- Actually: unravel into wavefunction stochastic dynamics: **Quantum State Diffusion**

T. Miura, Y. Akamatsu, M. Asakawa, S. Kajimoto, A.R., in preparation

Proof of
principle
in 1d



- Encouraging preliminary results: admixtures become independent of initial conditions
- Distribution of states at late times agrees well with Boltzmann and yields consistent T

Conclusion

- **Conceptual and technical progress** in in-medium quarkonium theory
- Recent and ongoing studies on quarkonium dynamical properties
 - Control over systematics in direct **spectral reconstructions** in lattice NRQCD
S.Kim, P. Petreczky, A.R., JHEP 1811 (2018) 088
 - **Pade based** extraction of the in-medium **heavy quark potential** possible
P. Petreczky, A.R., J. Weber, NPA982 (2019) 735
 - Improved **analytic parametrization** of $V(R)$ using the generalized **Gauss law**
D. Lafferty, A.R., arXiv:1906.00035
 - First consistent **Lindblad equation** for in-medium heavy quarkonium
with T. Miura, Y. Akamatsu, M. Asakawa (in progress)
- A lot of work remains to be done:
 - Explore the initial stages of a HIC: **formation dynamics** of quarkonium
with A. Lehmann (in preparation)
 - Improve reconstruction of spectral functions: **excited states physics**


Conclusion

- **Conceptual and technical progress** in in-medium quarkonium theory
- Recent and ongoing studies on quarkonium dynamical properties
 - Control over systematics in direct **spectral reconstructions** in lattice NRQCD
S.Kim, P. Petreczky, A.R., JHEP 1811 (2018) 088
 - **Pade based** extraction of the in-medium **heavy quark potential** possible
P. Petreczky, A.R., J. Weber, NPA982 (2019) 735
 - Improved **analytic parametrization** of $V(R)$ using the generalized **Gauss law**
D. Lafferty, A.R., arXiv:1906.00035
 - First consistent **Lindblad equation** for in-medium heavy quarkonium
with T. Miura, Y. Akamatsu, M. Asakawa (in progress)
- A lot of work remains to be done:
 - Explore the initial stages of a HIC: **formation dynamics** of quarkonium
with A. Lehmann (in preparation)
 - Improve reconstruction of spectral functions: **excited states physics**

Thank you for your attention


Extracting the potential

Example: Pade based reconstructions at $\beta=7.825$ $T=407\text{MeV}$ $N_t=12$
P. Petreczky, A.R. J. Weber in preparation




Extracting the potential

Example: Pade based reconstructions at $\beta=7.825$ $T=407\text{MeV}$ $N_T=12$
 P. Petreczky, A.R. J. Weber in preparation





Extracting the potential


Example: Pade based reconstructions at $\beta=7.825$ $T=407\text{MeV}$ $N_\tau=12$
 P. Petreczky, A.R. J. Weber in preparation



DFT




$$\rho(\omega) \approx -\frac{1}{\pi} \text{Im}[R_{N_\tau}(\omega)]$$




Extracting the potential

Example: Pade based reconstructions at $\beta=7.825$ $T=407\text{MeV}$ $N_\tau=12$
 P. Petreczky, A.R. J. Weber in preparation



- Always find well defined lowest peak: potential picture appears viable
- Beware of Pade artifacts besides peak: e.g. positivity violation, spikes



Stochastic potential

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

Stochastic potential

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

Stochastic potential

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- L, L^\dagger can be expressed in terms of $Re[V]$ and $Im[V]$ but e.o.m cannot be unravelled into a simple deterministic Schrödinger equation for ψ_{QQ}

Stochastic potential

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- L, L^\dagger can be expressed in terms of $Re[V]$ and $Im[V]$ but e.o.m cannot be unravelled into a simple deterministic Schrödinger equation for $\psi_{Q\bar{Q}}$
- Naïve Schroedinger vs. 1st order gradient exp.

$$\psi_{Q\bar{Q}}(t) = \exp \left[\underbrace{-\frac{\nabla^2}{M} + Re[V]}_{S_{pot}} + \underbrace{\eta(t)}_{S_{fluct}} \right] \psi_{Q\bar{Q}}(0)$$

Stochastic potential

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- L, L^\dagger can be expressed in terms of $Re[V]$ and $Im[V]$ but e.o.m cannot be unravelled into a simple deterministic Schrödinger equation for $\psi_{Q\bar{Q}}$
- Naïve Schroedinger vs. 1st order gradient exp.

$$\psi_{Q\bar{Q}}(t) = \exp \left[\underbrace{-\frac{\nabla^2}{M} + Re[V]}_{S_{pot}} + \underbrace{\eta(t)}_{S_{fluct}} \right] \psi_{Q\bar{Q}}(0)$$

$$i\partial_t \langle \psi_{Q\bar{Q}}(t) \rangle = \left(-\frac{\nabla^2}{M} + Re[V] - i|Im[V]| \right) \langle \psi_{Q\bar{Q}}(t) \rangle$$

Stochastic potential

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$


- L, L^\dagger can be expressed in terms of $Re[V]$ and $Im[V]$ but e.o.m cannot be unravelled into a simple deterministic Schrödinger equation for $\psi_{Q\bar{Q}}$

- Naïve Schroedinger vs. 1st order gradient exp.

$$\psi_{Q\bar{Q}}(t) = \exp \left[\underbrace{-\frac{\nabla^2}{M} + Re[V]}_{S_{pot}} + \underbrace{\eta(t)}_{S_{fluct}} \right] \psi_{Q\bar{Q}}(0)$$

$$i\partial_t \langle \psi_{Q\bar{Q}}(t) \rangle = \left(-\frac{\nabla^2}{M} + Re[V] - i|Im[V]| \right) \langle \psi_{Q\bar{Q}}(t) \rangle$$

GS survival probability



S.Kajimoto, Y.Akamatsu, M. Asakawa, A.R., PRD97 (2018) 014003

Stochastic potential

- Use scale separation: $m_Q > T$ heavy mass & weak coupling approximation

$$S_{FV} \approx S_{pot}[Re[V]] + S_{fluct}[Im[V]] + S_{diss}[Im[V]] + S_{LB}$$

- In QM language corresponds to Markovian evolution by Lindblad equation

$$\frac{d}{dt}\rho_{Q\bar{Q}}(t) = -i[H_{Q\bar{Q}}, \rho_{Q\bar{Q}}] + \sum_{i=1}^{N_{LB}} \gamma_i \left(\hat{L}_i \rho_{Q\bar{Q}} \hat{L}_i^\dagger - \frac{1}{2} \hat{L}_i \hat{L}_i^\dagger \rho_{Q\bar{Q}} - \frac{1}{2} \rho_{Q\bar{Q}} \hat{L}_i \hat{L}_i^\dagger \right)$$

- L, L^\dagger can be expressed in terms of $Re[V]$ and $Im[V]$ but e.o.m cannot be unravelled into a simple deterministic Schrödinger equation for $\psi_{Q\bar{Q}}$


- Naïve Schroedinger vs. 1st order gradient exp.

$$\psi_{Q\bar{Q}}(t) = \exp \left[\underbrace{-\frac{\nabla^2}{M} + Re[V]}_{S_{pot}} + \underbrace{\eta(t)}_{S_{fluct}} \right] \psi_{Q\bar{Q}}(0)$$

$$i\partial_t \langle \psi_{Q\bar{Q}}(t) \rangle = \left(-\frac{\nabla^2}{M} + Re[V] - i|Im[V]| \right) \langle \psi_{Q\bar{Q}}(t) \rangle$$

- Applicable at early times but incapable of thermalizing the heavy quark pair.

GS survival probability



S.Kajimoto, Y.Akamatsu, M. Asakawa, A.R., PRD97 (2018) 014003